

The expression of erythropoietin and its receptor in the developing rat retina

D Végvári^{1*}, A Szabó¹, G Deák², Á Lukács¹, V Doma¹, Ál Berta¹, Á Szél¹

¹Department of Human Morphology and Developmental Biology, Semmelweis University, Budapest, Hungary,

²Eye Clinic, Semmelweis University, Budapest, Hungary

Evidence from a number of studies indicates that the erythropoietin (EPO) does not only affect the haematopoietic system, but it exhibits neuroprotective and neurotrophic properties in the central nervous system and in the retina. Both the erythropoietin and its receptor (EPOR) are expressed in the retina, however, their cellular distribution and physiological role are not known. In the present study, we examined developing postnatal and adult rat retinas by EPO- and EPOR-specific antibodies with single and double labeling techniques in order to identify the temporal and spatial onset of their expression in the mammalian retina.

Retinas from Sprague-Dawley rats of different developmental ages were collected and analysed by immunohistochemistry. Following fixation and cryoprotection 10 µm thick frozen sections were cut from each specimen and stained with EPO and EPOR-specific antibodies. In double labelling experiments the horizontal cell-specific anti-calbindin antibody and the cone-specific marker PNA were also used. The labelling was visualised by the ABC method or by fluorescent antibodies and analysed by light or confocal microscopy.

At the time of birth the EPO-specific staining resulted a strong cytoplasmatic labelling in the ganglion cell layer and a weak, diffuse staining in the neuroblast layer. On the following days strong membrane immunoreactivity appeared in the developing horizontal and amacrine cells, and by the end of the first postnatal week the entire inner nuclear layer became positively stained. From the beginning of the second postnatal week on immunopositivity in the inner and outer plexiform layers and in the developing photoreceptor processes was also observed. The same result was found in the adult, where the cells of the ganglion cell layer and the inner nuclear layer, both plexiform layers and the photoreceptor inner segments bound the EPO-specific antibody. The staining pattern of the EPOR-specific antibody and the time of onset of the immunopositive elements showed big similarity to that of the EPO-specific antibody.

Since the spatial and temporal onset of the expression of the erythropoietin and its receptor correlates with the postnatal developmental events of rat retina, our findings suggest the involvement of the erythropoietin in the regulation of cell maturation, developmental apoptosis and synaptogenesis.

OTKA#T-042524, OTKA#F-61717.

*Corresponding author

E-mail: vegvári_dora@freemail.hu

Centrifugal visual fibers arising from the limbic system and the hypothalamus

V Vereczki^{1*}, Á Csáki², GE Hoffman³, G Fiskum⁴, K Kovács²

¹Department of Anatomy, Histology and Embryology, Semmelweis University of Medicine, Budapest, Hungary, ²Department of Human Morphology and Developmental Biology, Semmelweis University of Medicine, Budapest, Hungary, ³Department of Anatomy and Neurobiology, The University of Maryland School of Medicine, Baltimore, MD, USA, ⁴Department of Anesthesiology, The University of Maryland School of Medicine, Baltimore, MD, USA

It is already known that the optic nerve contains centrifugal fibers which are positive to VIP, LHRH or PACAP, but their origin had not yet been fully elucidated.

In our experiments utilizing the retrograde and anterograde transport of tracers (biotinylated dextran amine[BDA] and Fluorogold [FG], respectively) we have provided direct evidence for the cells of origin of a limboretinal and hypothalamoretinal pathways in rats and the termination of the centrifugal visual fibers in the retina using light microscopic approach. With the use of adult male rats following intravitreal injection of 10 000 kD BDA into the right eye, we have found retrogradely labeled